

# **Product datasheet for TA502998S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PDXK Mouse Monoclonal Antibody [Clone ID: OTI5H5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5H5

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDXK(NP\_03672) produced in HEK293 cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.82 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 34.9 kDa

**Gene Name:** pyridoxal kinase

Database Link: NP 003672

Entrez Gene 83578 RatEntrez Gene 216134 MouseEntrez Gene 8566 Human

000764

**Background:** The protein encoded by this gene phosphorylates vitamin B6, a step required for the

conversion of vitamin B6 to pyridoxal-5-phosphate, an important cofactor in intermediary

metabolism. The encoded protein is cytoplasmic and probably acts as a homodimer.

Alternatively spliced transcript variants have been described, but their biological validity has

not been determined. [provided by RefSeq, Jul 2008]

Synonyms: C21orf97; C21orf124; HEL-S-1a; PKH; PNK; PRED79







**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, Vitamin B6 metabolism

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PDXK (Cat# [RC200975], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDXK(Cat# [TA502998]). Positive lysates [LY418499] (100ug) and [LC418499] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 7 different cell lines by using anti-PDXK monoclonal antibody (1:200).






Equivalent amounts of cell lysates (10 ug per lane) of wild-type HEK293T cells (WT, Cat# LC810293T) and PDXK-Knockout HEK293T cells (KO, Cat# [LC841087]) were separated by SDS-PAGE and immunoblotted with anti-PDXK monoclonal antibody [TA502998] (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])






Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])






Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-PDXK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502998])



Anti-PDXK mouse monoclonal antibody ([TA502998]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PDXK ([RC200975]).



HEK293T cells transfected with either [RC200975] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDXK antibody ([TA502998]), and then analyzed by flow cytometry.