

### Product datasheet for TA503055BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SOCS3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3B8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3B8

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SOCS3 (NP\_003946) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 24.6 kDa

**Gene Name:** suppressor of cytokine signaling 3

Database Link: NP 003946

Entrez Gene 12702 MouseEntrez Gene 89829 RatEntrez Gene 9021 Human

014543





Background: This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as

suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene is induced by various cytokines, including IL6, IL10, and interferon (IFN)-gamma. The protein encoded by this gene can bind to JAK2 kinase, and inhibit the activity of JAK2 kinase. Studies of the mouse counterpart of this gene suggested the roles of this gene in the negative regulation of fetal

liver hematopoiesis, and placental development. [provided by RefSeq]


Synonyms: ATOD4; CIS3; Cish3; SOCS-3; SSI-3; SSI3

**Protein Families:** Druggable Genome


**Protein Pathways:** Adipocytokine signaling pathway, Insulin signaling pathway, Jak-STAT signaling pathway, Type

II diabetes mellitus, Ubiquitin mediated proteolysis

# **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SOCS3 ([RC209305], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SOCS3.




Anti-SOCS3 mouse monoclonal antibody ([TA503055]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SOCS3 ([RC209305]).









Flow cytometric Analysis of Jurkat cells, using anti-SOCS3 antibody ([TA503055]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).