

## Product datasheet for **TA503063M**

### **XPF (ERCC4) Mouse Monoclonal Antibody [Clone ID: OTI2D4]**

#### **Product data:**


|                         |  |
|-------------------------|--|
| Product Type:           | Primary Antibodies   |
| Clone Name:             | OTI2D4   |
| Applications:           | FC, IF, WB   |
| Recommended Dilution:   | WB 1:2000, IF 1:100, FLOW 1:100  |
| Reactivity:             | Human, Mouse   |
| Host:                   | Mouse  |
| Isotype:                | IgG2b  |
| Clonality:              | Monoclonal   |
| Immunogen:              | Full length human recombinant protein of human ERCC4 (NP_005227) produced in HEK293T cell.   |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.   |
| Concentration:          | 0.77 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)  |
| Conjugation:            | Unconjugated   |
| Storage:                | Store at -20°C as received.  |
| Stability:              | Stable for 12 months from date of receipt.   |
| Predicted Protein Size: | 104.3 kDa  |
| Gene Name:              | ERCC excision repair 4, endonuclease catalytic subunit   |
| Database Link:          | <a href="#">NP_005227</a><br><a href="#">Entrez Gene 50505 Mouse</a> <a href="#">Entrez Gene 2072 Human</a><br><a href="#">Q92889</a>  |
| Background:             | The protein encoded by this gene forms a complex with ERCC1 and is involved in the 5' incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that interacts with EME1. Defects in this gene are a cause of xeroderma pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6). |
| Synonyms:               | ERCC11; FANCC; RAD1; XFEPS; XPF  |


[View online »](#)


**Protein Families:** Druggable Genome

**Protein Pathways:** Nucleotide excision repair


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ERCC4 ([RC223300], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ERCC4. Positive lysates [LY401605] (100ug) and [LC401605] (20ug) can be purchased separately from OriGene.



Anti-ERCC4 mouse monoclonal antibody ([TA503063]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ERCC4 ([RC223300]).



HEK293T cells transfected with either [RC223300] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ERCC4 antibody ([TA503063]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-ERCC4 antibody ([TA503063M]), (Red), compared to a nonspecific negative control antibody, (Blue).