

Product datasheet for **TA503225S**

CBR3 Mouse Monoclonal Antibody [Clone ID: OTI1G6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1G6
Applications:	IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:150
Reactivity:	Human, Dog, Rat, Monkey, Mouse
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CBR3(NP_001227) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.43 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	30.7 kDa
Gene Name:	carbonyl reductase 3
Database Link:	NP_001227 Entrez Gene 109857 Mouse Entrez Gene 304078 Rat Entrez Gene 487748 Dog Entrez Gene 695769 Monkey Entrez Gene 874 Human O75828
Background:	Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq]




[View online »](#)

Synonyms: hCBR3; HEL-S-25; SDR21C2


Protein Families: Druggable Genome

Protein Pathways: Arachidonic acid metabolism, Metabolic pathways


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CBR3 (Cat# [RC201073], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CBR3(Cat# [TA503225]). Positive lysates [LY420049] (100ug) and [LC420049] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CBR3 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-CBR3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503225])