

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA503233BM

## SIRT5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2C4]

## **Product data:**

Product Type:	Primary Antibodies		
Clone Name:	OTI2C4		
Applications:	FC, IF, WB		
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100		
Reactivity:	Human, Mouse, Rat		
Host:	Mouse		
lsotype:	lgG1		
Clonality:	Monoclonal		
Immunogen:	Full length human recombinant protein of human SIRT5(NP_036373) produced in HEK293T cell.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.		
Concentration:	0.5 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	HRP		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	33.7 kDa		
Gene Name:	sirtuin 5		
Database Link:	<u>NP_036373</u> <u>Entrez Gene 68346 MouseEntrez Gene 306840 RatEntrez Gene 23408 Human</u> <u>Q9NXA8</u>		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### SIRT5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2C4] – TA503233BM

Background:This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2<br/>protein. Members of the sirtuin family are characterized by a sirtuin core domain and<br/>grouped into four classes. The functions of human sirtuins have not yet been determined;<br/>however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress<br/>recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular<br/>regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this<br/>gene is included in class III of the sirtuin family. Alternative splicing of this gene results in<br/>multiple transcript variants. [provided by RefSeq]

Synonyms:	SIR2L5


Protein Families:

#### Druggable Genome, Transcription Factors

## **Product images:**


170	-	
130	-	
100	-	
70	-	
55	-	
40	-	
35	-	
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SIRT5 ([RC200189], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SIRT5. Positive lysates [LY415885] (100ug) and [LC415885] (20ug) can be purchased separately from OriGene.



Anti-SIRT5 mouse monoclonal antibody ([TA503233]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SIRT5 ([RC200189]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



HEK293T cells transfected with either [RC200189] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SIRT5 antibody ([TA503233]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US