

# Product datasheet for TA503240AM

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **OBFC1 (STN1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2E4]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2E4

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human OBFC1 (NP\_079204) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 41.9 kDa

**Gene Name:** STN1 subunit of CST complex

Database Link: NP 079204

Entrez Gene 79991 Human

Q9H668

Background: OBFC1 and C17ORF68 (MIM 613129) are subunits of an alpha accessory factor (AAF) that

stimulates the activity of DNA polymerase-alpha-primase (see MIM 176636), the enzyme that initiates DNA replication (Casteel et al., 2009 [PubMed 19119139]). OBFC1 also appears to function in a telomere-associated complex with C17ORF68 and TEN1 (C17ORF106; MIM

613130) (Miyake et al., 2009 [PubMed 19854130]). [supplied by OMIM]







Synonyms:

AAF-44; AAF44; bA541N10.2; RPA-32; STN1


# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY OBFC1 (Cat# [RC200778], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OBFC1(Cat# [TA503240]). Positive lysates [LY410967] (100ug) and [LC410967] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-OBFC1 monoclonal antibody at 1:200 dilution.



Anti-OBFC1 mouse monoclonal antibody ([TA503240]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY OBFC1 ([RC200778]).



HEK293T cells transfected with either [RC200778] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-OBFC1 antibody ([TA503240]), and then analyzed by flow cytometry.