

### **Product datasheet for TA503283**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### SIRT5 Mouse Monoclonal Antibody [Clone ID: OTI2E10]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2E10
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SIRT5 (NP\_036373) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.61 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 33.7 kDa

Gene Name: sirtuin 5

Database Link: NP 036373

Entrez Gene 68346 MouseEntrez Gene 306840 RatEntrez Gene 23408 Human

O9NXA8






#### Background:

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class III of the sirtuin family. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq]


Synonyms: SIR2L5

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SIRT5 ([RC200189], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SIRT5. Positive lysates [LY415885] (100ug) and [LC415885] (20ug) can be purchased separately from OriGene.



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and SIRT5-Knockout HeLa cells (KO, Cat# [LC830483]) were separated by SDS-PAGE and immunoblotted with anti-SIRT5 monoclonal antibody TA503283 (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.





HEK293T cells transfected with either [RC200189] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SIRT5 antibody (TA503283), and then analyzed by flow cytometry.