

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA503336M

PDE1B Mouse Monoclonal Antibody [Clone ID: OTI1D6]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI1D6	
Applications:	IF, WB	
Recommended Dilution:	WB 1:2000, IF 1:100	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human PDE1B(NP_000915) produced in HEK293T cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	0.74 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	61.2 kDa	
Gene Name:	phosphodiesterase 1B	
Database Link:	<u>NP_000915</u> <u>Entrez Gene 18574 MouseEntrez Gene 29691 RatEntrez Gene 5153 Human</u> <u>Q01064</u>	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PDE1B Mouse Monoclonal Antibody [Clone ID: OTI1D6] – TA503336M
Background:	The protein encoded by this gene belongs to the cyclic nucleotide phosphodiesterase (PDE) family, and PDE1 subfamily. Members of the PDE1 family are calmodulin-dependent PDEs that are stimulated by a calcium-calmodulin complex. This PDE has dual-specificity for the second messengers, cAMP and cGMP, with a preference for cGMP as a substrate. cAMP and cGMP function as key regulators of many important physiological processes. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2011]
Synonyms:	PDE1B1; PDES1B
Protein Families:	Druggable Genome
Protein Pathways	Calcium signaling pathway, Progesterone-mediated oocyte maturation, Purine metabolism

Product images:

170	_	
130	-	
100	-	
70	-	- 10
55	-	
40	-	
35	-	
25	-	
15	-1	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDE1B ([RC206588], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDE1B. Positive lysates [LY400337] (100ug) and [LC400337] (20ug) can be purchased separately from OriGene.



Anti-PDE1B mouse monoclonal antibody ([TA503336]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PDE1B ([RC206588]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US