

Product datasheet for TA503340S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBOX5 Mouse Monoclonal Antibody [Clone ID: OTI1C5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C5

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:200~500, IHC 1:150, FLOW 1:100, IF: 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-130 and C 419-487 of

human UBOX5(NP_955447) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: U-box domain containing 5

Database Link: NP 955447

Entrez Gene 22888 Human

<u>094941</u>

Synonyms: hUIP5; RNF37; UBCE7IP5; UIP5


Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY UBOX5 (Cat# [RC208505], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UBOX5(Cat# [TA503340]).




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-UBOX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503340])




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-UBOX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503340])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-UBOX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503340])




Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-UBOX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503340])




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-UBOX5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503340])






Anti-UBOX5 mouse monoclonal antibody ([TA503340]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UBOX5 ([RC208505]).




HEK293T cells transfected with either [RC208505] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-UBOX5 antibody ([TA503340]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-UBOX5 antibody ([TA503340]), (Red), compared to a nonspecific negative control antibody, (Blue).





Flow cytometric Analysis of Jurkat cells, using anti-UBOX5 antibody ([TA503340]), (Red), compared to a nonspecific negative control antibody, (Blue).