

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA503415

Cytochrome P450 17A1 (CYP17A1) Mouse Monoclonal Antibody [Clone ID: OTI5G10]

Product data:

Product Type:	Primary Antibodies		
Clone Name:	OTI5G10		
Applications:	WB		
Recommended Dilution:	WB 1:2000		
Reactivity:	Human		
Host:	Mouse		
lsotype:	lgG1		
Clonality:	Monoclonal		
Immunogen:	Full length human recombinant protein of human CYP17A1(NP_000093) produced in HEK293T cell.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Concentration:	1 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	Unconjugated		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	57.2 kDa		
Gene Name:	cytochrome P450 family 17 subfamily A member 1		
Database Link:	<u>NP_000093</u> <u>Entrez Gene 1586 Human</u> <u>P05093</u>		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Cytochrome P450 17A1 (CYP17A1) Mouse Monoclonal Antibody [Clone ID: OTI5G10] – TA503415
Background:	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. It has both 17alpha-hydroxylase and 17,20-lyase activities and is a key enzyme in the steroidogenic pathway that produces progestins, mineralocorticoids, glucocorticoids, androgens, and estrogens. Mutations in this gene are associated with isolated steroid-17 alpha-hydroxylase deficiency, 17-alpha-hydroxylase/17,20-lyase deficiency, pseudohermaphroditism, and adrenal hyperplasia. [provided by RefSeq]
Synonyms:	CPT7; CYP17; P450C17; S17AH
Protein Families:	Druggable Genome, P450
Protein Pathway	s: C21-Steroid hormone metabolism, Metabolic pathways

Product images:

170	_	
130	-	
100	_	
70	_	
55	-	-
40	_	
35	—	
25	-	
15	_	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CYP17A1 ([RC209042], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CYP17A1. Positive lysates [LY400030] (100ug) and [LC400030] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US