

Product datasheet for **TA503506S**

PNPO Mouse Monoclonal Antibody [Clone ID: OTI1G9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1G9
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:4000, IHC 1:150, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PNPO(NP_060599) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.8 kDa
Gene Name:	pyridoxamine 5'-phosphate oxidase
Database Link:	NP_060599 Entrez Gene 64533 Rat Entrez Gene 103711 Mouse Entrez Gene 55163 Human Q9NVS9
Background:	The enzyme encoded by this gene catalyzes the terminal, rate-limiting step in the synthesis of pyridoxal 5'-phosphate, also known as vitamin B6. Vitamin B6 is a required co-factor for enzymes involved in both homocysteine metabolism and synthesis of neurotransmitters such as catecholamine. Mutations in this gene result in pyridoxamine 5'-phosphate oxidase (PNPO) deficiency, a form of neonatal epileptic encephalopathy. [provided by RefSeq]. COMPLETENESS: complete on the 3' end.




[View online »](#)


Synonyms: HEL-S-302; PDXPO

Protein Pathways: Metabolic pathways, Vitamin B6 metabolism


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PNPO (Cat# [RC200133], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PNPO (Cat# [TA503506]). Positive lysates [LY413282] (100ug) and [LC413282] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-PNPO mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503506])



Flow cytometric Analysis of HeLa cells, using anti-PNPO antibody ([TA503506]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-PNPO antibody ([TA503506]), (Red), compared to a nonspecific negative control antibody, (Blue).