

## Product datasheet for **TA503534**

### UNG Mouse Monoclonal Antibody [Clone ID: OTI2F8]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2F8
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human UNG(NP_550433) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.57 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	34.5 kDa
Gene Name:	uracil DNA glycosylase
Database Link:	<a href="#">NP_550433</a> <a href="#">Entrez Gene 22256 Mouse</a> <a href="#">Entrez Gene 304577 Rat</a> <a href="#">Entrez Gene 7374 Human</a> <a href="#">P13051</a>



[View online »](#)

**Background:**

This gene encodes one of several uracil-DNA glycosylases. One important function of uracil-DNA glycosylases is to prevent mutagenesis by eliminating uracil from DNA molecules by cleaving the N-glycosylic bond and initiating the base-excision repair (BER) pathway. Uracil bases occur from cytosine deamination or misincorporation of dUMP residues. Alternative promoter usage and splicing of this gene leads to two different isoforms: the mitochondrial UNG1 and the nuclear UNG2. The UNG2 term was used as a previous symbol for the CCNO gene (GenelD 10309), which has been confused with this gene, in the literature and some databases. [provided by RefSeq]

**Synonyms:**


DGU; HIGM4; HIGM5; UDG; UNG1; UNG2; UNG15

**Protein Families:**


Druggable Genome, Stem cell - Pluripotency

**Protein Pathways:**


Base excision repair, Primary immunodeficiency

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UNG ([RC222868], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UNG. Positive lysates [LY408998] (100ug) and [LC408998] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-UNG mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503534)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-UNG mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503534)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-UNG mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503534)



Anti-UNG mouse monoclonal antibody (TA503534) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UNG ([RC222868]).