

# Product datasheet for TA503567S

# UNG Mouse Monoclonal Antibody [Clone ID: OTI2F3]

# **Product data:**

#### **Product Type: Primary Antibodies Clone Name:** OTI2F3 **Applications:** IF, WB Recommended Dilution: WB 1:2000, IF 1:100 **Reactivity:** Human, Mouse, Rat Host: Mouse Isotype: lgG2a **Clonality:** Monoclonal Full length human recombinant protein of human UNG(NP\_550433) produced in HEK293T Immunogen: cell. Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. Concentration: 1.1 mg/ml **Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) **Conjugation:** Unconjugated Store at -20°C as received. Storage: Stability: Stable for 12 months from date of receipt. Predicted Protein Size: 34.5 kDa Gene Name: uracil DNA glycosylase Database Link: NP 550433 Entrez Gene 22256 MouseEntrez Gene 304577 RatEntrez Gene 7374 Human P13051



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


### **GRIGENE** UNG Mouse Monoclonal Antibody [Clone ID: OTI2F3] – TA503567S

- Background: This gene encodes one of several uracil-DNA glycosylases. One important function of uracil-DNA glycosylases is to prevent mutagenesis by eliminating uracil from DNA molecules by cleaving the N-glycosylic bond and initiating the base-excision repair (BER) pathway. Uracil bases occur from cytosine deamination or misincorporation of dUMP residues. Alternative promoter usage and splicing of this gene leads to two different isoforms: the mitochondrial UNG1 and the nuclear UNG2. The UNG2 term was used as a previous symbol for the CCNO gene (GeneID 10309), which has been confused with this gene, in the literature and some databases. [provided by RefSeq]
  Synonyms: DGU; HIGM4; HIGM5; UDG; UNG1; UNG2; UNG15
- Protein Families: Druggable Genome, Stem cell Pluripotency
- Protein Pathways: Base excision repair, Primary immunodeficiency

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UNG ([RC222868], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UNG. Positive lysates [LY408998] (100ug) and [LC408998] (20ug) can be purchased separately from OriGene.



Anti-UNG mouse monoclonal antibody ([TA503567]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UNG ([RC222868]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US