

Product datasheet for TA503572AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

AIPL1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1B2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1B2

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human AIPL1(NP_055151) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.7 kDa

Gene Name: aryl hydrocarbon receptor interacting protein like 1

Database Link: NP 055151

Entrez Gene 59110 RatEntrez Gene 114230 MouseEntrez Gene 23746 Human

O9NZN9






Background:

Leber congenital amaurosis (LCA) accounts for at least 5% of all inherited retinal disease and is the most severe inherited retinopathy with the earliest age of onset. Individuals affected with LCA are diagnosed at birth or in the first few months of life with severely impaired vision or blindness, nystagmus and an abnormal or flat electroretinogram. The photoreceptor/pineal -expressed gene, AIPL1, encoding aryl-hydrocarbon interacting protein-like 1, was mapped within the LCA4 candidate region. The protein contains three tetratricopeptide motifs, consistent with nuclear transport or chaperone activity. AIPL1 mutations may cause approximately 20% of recessive LCA. [provided by RefSeq, Jul 2008]


Synonyms: AIPL2; LCA4

Protein Families: Druggable Genome


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY AIPL1 ([RC204079], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AIPL1. Positive lysates [LY415353] (100ug) and [LC415353] (20ug) can be purchased separately from OriGene.



Anti-AIPL1 mouse monoclonal antibody ([TA503572]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY AIPL1 ([RC204079]).



HEK293T cells transfected with either [RC204079] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-AIPL1 antibody ([TA503572]), and then analyzed by flow cytometry.