

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA503632BM

Acetyl CoA synthetase (ACSS2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3C6]

Product data:

Product Type:	Primary Antibodies		
Clone Name:	OTI3C6		
Applications:	WB		
Recommended Dilution:	WB 1:2000		
Reactivity:	Human, Mouse, Rat		
Host:	Mouse		
lsotype:	lgG2b		
Clonality:	Monoclonal		
Immunogen:	Full length human recombinant protein of human ACSS2(NP_061147) produced in HEK293T cell.		
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.		
Concentration:	0.5 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Conjugation:	HRP		
Storage:	Store at -20°C as received.		
Stability:	Stable for 12 months from date of receipt.		
Predicted Protein Size:	78.4 kDa		
Gene Name:	acyl-CoA synthetase short chain family member 2		
Database Link:	<u>NP_061147</u> <u>Entrez Gene 60525 MouseEntrez Gene 311569 RatEntrez Gene 55902 Human</u> <u>Q9NR19</u>		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Acetyl CoA synthetase (ACSS2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3C6] – TA503632BM
Background:	This gene encodes a cytosolic enzyme that catalyzes the activation of acetate for use in lipid synthesis and energy generation. The protein acts as a monomer and produces acetyl-CoA from acetate in a reaction that requires ATP. Expression of this gene is regulated by sterol regulatory element-binding proteins, transcription factors that activate genes required for the synthesis of cholesterol and unsaturated fatty acids. Alternative splicing results in multiple transcript variants. [provided by RefSeq]
Synonyms:	ACAS2; ACECS; ACS; ACSA; dJ1161H23.1
Protein Pathway	s: Glycolysis / Gluconeogenesis, Metabolic pathways, Propanoate metabolism, Pyruvate metabolism

Product images:

		1000	
170	_		
130	-		
100	_		
70	_		
55	_		
40	—		
35	—		
25	-		
15	-		
10	-		

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACSS2 ([RC204260], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACSS2. Positive lysates [LY412981] (100ug) and [LC412981] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US