

Product datasheet for TA503684AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Acetyl CoA synthetase (ACSS2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2E9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2E9

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ACSS2(NP_061147) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 78.4 kDa

Gene Name: acyl-CoA synthetase short chain family member 2

Database Link: NP 061147

Entrez Gene 60525 MouseEntrez Gene 311569 RatEntrez Gene 55902 Human

Q9NR19





Acetyl CoA synthetase (ACSS2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2E9] – TA503684AM

Background: This gene encodes a cytosolic enzyme that catalyzes the activation of acetate for use in lipid

synthesis and energy generation. The protein acts as a monomer and produces acetyl-CoA from acetate in a reaction that requires ATP. Expression of this gene is regulated by sterol regulatory element-binding proteins, transcription factors that activate genes required for the synthesis of cholesterol and unsaturated fatty acids. Alternative splicing results in multiple


transcript variants. [provided by RefSeq]

Synonyms: ACAS2; ACECS; ACS, ACSA; dJ1161H23.1

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, Propanoate metabolism, Pyruvate

metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACSS2 ([RC204260], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACSS2. Positive lysates [LY412981] (100ug) and [LC412981] (20ug) can be purchased separately from OriGene.