

Product datasheet for TA503770AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2M Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2D9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2D9

Applications: IF, WB

Recommended Dilution: WB 1:500~2000, IF 1:100

Reactivity: Human, Dog, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human UBE2M(NP_003960) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 20.7 kDa

Gene Name: ubiquitin conjugating enzyme E2 M

Database Link: NP 003960

Entrez Gene 22192 MouseEntrez Gene 361509 RatEntrez Gene 484221 DogEntrez Gene

712206 MonkeyEntrez Gene 9040 Human

P61081






Background:

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is linked with a ubiquitin-like protein, NEDD8, which can be conjugated to cellular proteins, such as Cdc53/culin. [provided by RefSeq]


Synonyms: hUbc12; UBC-RS2; UBC12

Protein Pathways: Ubiquitin mediated proteolysis

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY UBE2M (Cat# [RC208946], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UBE2M(Cat# [TA503770]). Positive lysates [LY418316] (100ug) and [LC418316] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-UBE2M monoclonal antibody.





Anti-UBE2M mouse monoclonal antibody ([TA503770]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UBE2M ([RC208946]).