

Product datasheet for TA503784AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Lipoprotein lipase (LPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6B4

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding amino acids 28-475 of human

LPL(NP_000228) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50.3 kDa

Gene Name: lipoprotein lipase

Database Link: NP 000228

Entrez Gene 16956 MouseEntrez Gene 24539 RatEntrez Gene 4023 Human

P06858

Background: LPL encodes lipoprotein lipase, which is expressed in heart, muscle, and adipose tissue. LPL

functions as a homodimer, and has the dual functions of triglyceride hydrolase and

ligand/bridging factor for receptor-mediated lipoprotein uptake. Severe mutations that cause LPL deficiency result in type I hyperlipoproteinemia, while less extreme mutations in LPL are

linked to many disorders of lipoprotein metabolism. [provided by RefSeq]






Lipoprotein lipase (LPL) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6B4] – TA503784AM

Synonyms: HDLCQ11; LIPD

Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Glycerolipid metabolism, PPAR signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY LPL ([RC203766], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-LPL. Positive lysates [LY400089] (100ug) and [LC400089] (20ug) can be purchased separately from OriGene.