

Product datasheet for TA503801S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Adiponectin (ADIPOQ) Mouse Monoclonal Antibody [Clone ID: OTI1B2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1B2

Applications: FC, IF, IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 19-244 of human

ADIPOQ(NP_004788) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 24.9 kDa

Gene Name: adiponectin, C1Q and collagen domain containing

Database Link: NP 004788

Entrez Gene 9370 Human

Q15848

Background: This gene is expressed in adipose tissue exclusively. It encodes a protein with similarity to

collagens X and VIII and complement factor C1q. The encoded protein circulates in the plasma and is involved with metabolic and hormonal processes. Mutations in this gene are associated with adiponectin deficiency. Multiple alternatively spliced variants, encoding the

same protein, have been identified. [provided by RefSeg, Apr 2010]





Synonyms: ACDC; ACRP30; ADIPQTL1; ADPN; APM-1; APM1; GBP28


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Adipocytokine signaling pathway, PPAR signaling pathway, Type II diabetes mellitus


Product images:









Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-ADIPOQ monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ADIPOQ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503801])




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-ADIPOQ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503801])




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-ADIPOQ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503801])





Anti-ADIPOQ mouse monoclonal antibody ([TA503801]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ADIPOQ ([RC215161]).



HEK293T cells transfected with either [RC215161] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ADIPOQ antibody ([TA503801]), and then analyzed by flow cytometry.