

### **Product datasheet for TA503824BM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Heme oxygenase 2 (HMOX2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4H6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4H6
Applications: FC, WB

**Reactivity:** WB 1:500, FLOW 1:100 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human HMOX2(NP\_002125) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 35.9 kDa

**Gene Name:** heme oxygenase 2

Database Link: NP 002125

Entrez Gene 15369 MouseEntrez Gene 79239 RatEntrez Gene 3163 Human

P30519



## Heme oxygenase 2 (HMOX2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4H6] – TA503824BM

Background:


Heme oxygenase, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is subsequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitter. Heme oxygenase activity is induced by its substrate heme and by various nonheme substances. Heme oxygenase occurs as 2 isozymes, an inducible heme oxygenase-1 and a constitutive heme oxygenase-2. HMOX1 and HMOX2 belong to the heme oxygenase family. Alternative splice variants encoding the same protein have been identified at this locus. [provided by RefSeq, Jul 2008]

Synonyms: HO-2


**Protein Families:** Transmembrane

**Protein Pathways:** Porphyrin and chlorophyll metabolism

#### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HMOX2 ([RC201777], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HMOX2. Positive lysates [LY419512] (100ug) and [LC419512] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC201777] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-HMOX2 antibody ([TA503824]), and then analyzed by flow cytometry.