

# Product datasheet for TA503833M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Acetyl CoA synthetase (ACSS2) Mouse Monoclonal Antibody [Clone ID: OTI2G8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G8
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human ACSS2(NP\_061147) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.43 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 78.4 kDa

**Gene Name:** acyl-CoA synthetase short chain family member 2

Database Link: NP 061147

Entrez Gene 60525 MouseEntrez Gene 311569 RatEntrez Gene 55902 Human

O9NR19

**Background:** This gene encodes a cytosolic enzyme that catalyzes the activation of acetate for use in lipid

synthesis and energy generation. The protein acts as a monomer and produces acetyl-CoA from acetate in a reaction that requires ATP. Expression of this gene is regulated by sterol regulatory element-binding proteins, transcription factors that activate genes required for the synthesis of cholesterol and unsaturated fatty acids. Alternative splicing results in multiple

transcript variants. [provided by RefSeq]




Synonyms: ACAS2; ACECS; ACSA; dJ1161H23.1


**Protein Pathways:** Glycolysis / Gluconeogenesis, Metabolic pathways, Propanoate metabolism, Pyruvate

metabolism

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACSS2 ([RC204260], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACSS2. Positive lysates [LY412981] (100ug) and [LC412981] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC204260] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ACSS2 antibody ([TA503833]), and then analyzed by flow cytometry.