

# **Product datasheet for TA503883S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ERp72 (PDIA4) Mouse Monoclonal Antibody [Clone ID: OTI2F1]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F1
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDIA4(NP\_004902) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.52 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 72.8 kDa

**Gene Name:** protein disulfide isomerase family A member 4

Database Link: NP 004902

Entrez Gene 12304 MouseEntrez Gene 116598 RatEntrez Gene 9601 Human

P13667

**Synonyms:** ERp-72; ERP70; ERP72


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Vibrio cholerae infection







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDIA4 ([RC204041], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDIA4. Positive lysates [LY401531] (100ug) and [LC401531] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PDIA4 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and PDIA4-Knockout 293T cells (KO, Cat# [LC841833]) were separated by SDS-PAGE and immunoblotted with anti-PDIA4 monoclonal antibody [TA503883], (1:500). Then the blotted membrane was stripped and reprobed with anti-b-actin antibody ([TA811000]) as a loading control.