

Product datasheet for TA503920BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CCK4 (PTK7) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2E7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2E7

Applications: FC, IF, WB

Recommended Dilution: WB 1:200~500, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PTK7(NP_690619) produced in HEK293T

cell

Formulation: PBS (pH 7.4) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: protein tyrosine kinase 7 (inactive)

Database Link: NP 690619

Entrez Gene 71461 MouseEntrez Gene 5754 Human

Q13308

Background: Receptor protein tyrosine kinases transduce extracellular signals across the cell membrane. A

subgroup of these kinases lack detectable catalytic tyrosine kinase activity but retain roles in signal transduction. The protein encoded by this gene is a member of this subgroup of tyrosine kinases and may function as a cell adhesion molecule. This gene is thought to be expressed in colon carcinomas but not in normal colon, and therefore may be a marker for or may be involved in tumor progression. Four transcript variants encoding four different

isoforms have been found for this gene. [provided by RefSeq, Jul 2008]







Synonyms: CCK-4; CCK4

Protein Families: Druggable Genome, Protein Kinase, Transmembrane


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PTK7 (Cat# [RC223416], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PTK7(Cat# [TA503920]).




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PTK7 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Anti-PTK7 mouse monoclonal antibody ([TA503920]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PTK7 ([RC223416]).





HEK293T cells transfected with either [RC223416] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PTK7 antibody ([TA503920]), and then analyzed by flow cytometry.