

## **Product datasheet for TA503936**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# PDSS2 Mouse Monoclonal Antibody [Clone ID: OTI1D10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1D10

Applications: FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDSS2(NP\_065114) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.8 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 43.9 kDa

**Gene Name:** decaprenyl diphosphate synthase subunit 2

Database Link: NP 065114

Entrez Gene 71365 MouseEntrez Gene 365592 RatEntrez Gene 57107 Human

086YH6

**Background:** The protein encoded by this gene is an enzyme that synthesizes the prenyl side-chain of

coenzyme Q, or ubiquinone, one of the key elements in the respiratory chain. The gene product catalyzes the formation of all trans-polyprenyl pyrophosphates from isopentyl diphosphate in the assembly of polyisoprenoid side chains, the first step in coenzyme Q biosynthesis. Defects in this gene are a cause of coenzyme Q10 deficiency. COMPLETENESS:

complete on the 3' end.







Synonyms: bA59I9.3; C6orf210; COQ10D3; DLP1; hDLP1

**Protein Pathways:** Terpenoid backbone biosynthesis


### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDSS2 ([RC207892], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDSS2. Positive lysates [LY412512] (100ug) and [LC412512] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503936)




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503936)







Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503936)



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503936)



Flow cytometric Analysis of Hela cells, using anti-PDSS2 antibody (TA503936), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-PDSS2 antibody (TA503936), (Red), compared to a nonspecific negative control antibody, (Blue).