

Product datasheet for TA503937

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DPP9 Mouse Monoclonal Antibody [Clone ID: OTI2F10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F10
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DPP9(NP_631898) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.67 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 96.4 kDa

Gene Name: dipeptidyl peptidase 9

Database Link: NP 631898

Entrez Gene 224897 MouseEntrez Gene 301130 RatEntrez Gene 91039 Human

O86TI2







Background:

This gene encodes a protein that is a member of the S9B family in clan SC of the serine proteases. The protein has been shown to have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. Although the activity of this protein is similar to that of dipeptidyl peptidase 4 (DPP4), it does not appear to be membrane bound. In general, dipeptidyl peptidases appear to be involved in the regulation of the activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. Several transcript variants of this gene have been described but not fully characterized. [provided by RefSeq]

Synonyms: DP9; DPLP9; DPRP-2; DPRP2
Protein Families: Druggable Genome, Protease

Product images:







HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DPP9 ([RC224465], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DPP9. Positive lysates [LY403380] (100ug) and [LC403380] (20ug) can be purchased separately from OriGene.

Figure from citation: Western blot analysis of DPP9 protein expression by using anti-DPP9 antibody in Huh7 cells were transfected with DPP9 siRNA or nontargeting (NT) control siRNA for 48 h, or treated with enzyme inhibitor, 8 uM 1G244 (or vehicle control, 0.08% DMSO) for 6 h before incubating on fibronectin-coated plastic for 30 min. Dilution: 1:2000 View Citation





Anti-DPP9 mouse monoclonal antibody (TA503937) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DPP9 ([RC224465]).



HEK293T cells transfected with either [RC224465] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DPP9 antibody (TA503937), and then analyzed by flow cytometry.