

Product datasheet for **TA503951AM**

PDSS2 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1D12]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D12
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:150, FLOW 1:100
Reactivity:	Human, Monkey, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDSS2(NP_065114) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	43.9 kDa
Gene Name:	decaprenyl diphosphate synthase subunit 2
Database Link:	NP_065114 Entrez Gene 71365 Mouse Entrez Gene 365592 Rat Entrez Gene 698357 Monkey Entrez Gene 57107 Human Q86YH6



[View online »](#)

Background:

The protein encoded by this gene is an enzyme that synthesizes the prenyl side-chain of coenzyme Q, or ubiquinone, one of the key elements in the respiratory chain. The gene product catalyzes the formation of all trans-polyprenyl pyrophosphates from isopentyl diphosphate in the assembly of polyisoprenoid side chains, the first step in coenzyme Q biosynthesis. Defects in this gene are a cause of coenzyme Q10 deficiency. **COMPLETENESS:** complete on the 3' end.


Synonyms:

bA59I9.3; C6orf210; COQ10D3; DLP1; hDLP1


Protein Pathways:

Terpenoid backbone biosynthesis


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PDSS2 (Cat# [RC207892], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDSS2(Cat# [TA503951]). Positive lysates [LY412512] (100ug) and [LC412512] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PDSS2 monoclonal antibody.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503951])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503951])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503951])



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-PDSS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503951])



HEK293T cells transfected with either [RC207892] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDSS2 antibody ([TA503951]), and then analyzed by flow cytometry.