

Product datasheet for **TA503968AM**

Kv beta 1 (KCNAB1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7C7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI7C7
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human KCNAB1(NP_751891) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	44.5 kDa
Gene Name:	potassium voltage-gated channel subfamily A regulatory beta subunit 1
Database Link:	NP_751891 Entrez Gene 16497 Mouse Entrez Gene 29737 Rat Entrez Gene 7881 Human Q14722



[View online »](#)

Background:

Potassium channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in *Drosophila*, and each has been shown to have human homolog(s). This gene encodes a member of the potassium channel, voltage-gated, shaker-related subfamily. This member includes three distinct isoforms which are encoded by three alternatively spliced transcript variants of this gene. These three isoforms are beta subunits, which form heteromultimeric complex with alpha subunits and modulate the activity of the pore-forming alpha subunits. [provided by RefSeq]


Synonyms:

AKR6A3; hKvb3; hKvBeta3; KCNA1B; KV-BETA-1; Kvb1.3


Protein Families:

Druggable Genome, Ion Channels: Other


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KCNA1 ([RC207384], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KCNA1. Positive lysates [LY403531] (100ug) and [LC403531] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-KCNA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503968])