

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA503992AM

## SDS Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI4E1]

### **Product data:**

| Product Type:           | Primary Antibodies  |
|-------------------------|---|
| Clone Name:             | OTI4E1  |
| Applications:           | FC, WB  |
| Recommended Dilution:   | WB 1:2000, FLOW 1:100   |
| Reactivity:             | Human   |
| Host:                   | Mouse   |
| lsotype:                | lgG1  |
| Clonality:              | Monoclonal  |
| Immunogen:              | Full length human recombinant protein of human SDS(NP_006834) produced in HEK293T<br>cell.  |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| Concentration:          | 0.5 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)  |
| Conjugation:            | Biotin  |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 34.4 kDa  |
| Gene Name:              | serine dehydratase  |
| Database Link:          | <u>NP_006834</u><br><u>Entrez Gene 10993 Human</u><br><u>P20132</u>   |
| Background:             | This gene encodes one of three enzymes that are involved in metabolizing serine and glycine.<br>L-serine dehydratase converts L-serine to pyruvate and ammonia and requires pyridoxal<br>phosphate as a cofactor. The encoded protein can also metabolize threonine to NH4+ and 2-<br>ketobutyrate. The encoded protein is found predominantly in the liver. [provided by RefSeq].<br>COMPLETENESS: complete on the 3' end. |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### Synonyms:


**Protein Pathways:** 

Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways

### **Product images:**



SDH



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SDS ([RC217814], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SDS. Positive lysates [LY416388] (100ug) and [LC416388] (20ug) can be purchased separately from OriGene.

HEK293T cells transfected with either [RC217814] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-SDS antibody ([TA503992]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US