

## **Product datasheet for TA504015AM**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### FBXO21 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6E7]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI6E7

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FBXO21(NP\_296373) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 72.1 kDa

**Gene Name:** F-box protein 21

Database Link: NP 296373

Entrez Gene 231670 MouseEntrez Gene 360818 RatEntrez Gene 23014 Human

094952






#### Background:


This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. Alternative splicing of this gene generates 2 transcript variants. [provided by RefSeq]

Synonyms: FBX21


# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FBXO21 ([RC223095], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FBXO21.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])






Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])







Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-FBXO21 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504015])



Anti-FBXO21 mouse monoclonal antibody ([TA504015]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY FBXO21 ([RC223095]).







Flow cytometric Analysis of Jurkat cells, using anti-FBXO21 antibody ([TA504015]), (Red), compared to a nonspecific negative control antibody, (Blue).