

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA504026AM

TPSG1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2H5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2H5
Applications:	WB
Recommended Dilution:	WB 1:200
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 20-283 of human TPSG1(NP_036599) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	32 kDa
Gene Name:	tryptase gamma 1
Database Link:	<u>NP_036599</u> <u>Entrez Gene 25823 Human</u> <u>Q9NRR2</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Serigene TPSG1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2H5] – TA504026AM

Background:Tryptases comprise a family of trypsin-like serine proteases, the peptidase family S1.
Tryptases are enzymatically active only as heparin-stabilized tetramers, and they are resistant
to all known endogenous proteinase inhibitors. Several tryptase genes are clustered on
chromosome 16p13.3. There is uncertainty regarding the number of genes in this cluster.
Currently four functional genes - alpha I, beta I, beta II and gamma I - have been identified.
And beta I has an allelic variant named alpha II, beta II has an allelic variant beta III, also
gamma I has an allelic variant gamma II. Beta tryptases appear to be the main isoenzymes
expressed in mast cells; whereas in basophils, alpha-tryptases predominant. This gene differs
from other members of the tryptase gene family in that it has C-terminal hydrophobic
domain, which may serve as a membrane anchor. Tryptases have been implicated as
mediators in the pathogenesis of asthma and other allergic and inflammatory disorders.
[provided by RefSeq, Jul 2008]. COMPLETENESS: complete on the 3' end.


Synonyms: PRSS31; TMT; trpA

Protein Families:

кэээт, тип, црА

milies: Druggable Genome, Transmembrane

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TPSG1 ([RC222359], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TPSG1. Positive lysates [LY402219] (100ug) and [LC402219] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US