

Product datasheet for TA504050AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DGKA Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI6A11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6A11
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DGKA(NP_001336) produced in HEK293

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 82.5 kDa

Gene Name: diacylglycerol kinase alpha

Database Link: NP 001336

Entrez Gene 1606 Human

P23743

Background: The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It

acts as a modulator that competes with protein kinase C for the second messenger diacylglycerol in intracellular signaling pathways. It also plays an important role in the resynthesis of phosphatidylinositols and phosphorylating diacylglycerol to phosphatidic acid.

Alternative splicing occurs at this locus and four transcript variants encoding the same

protein have been identified. [provided by RefSeq]






Synonyms: DAGK; DAGK1; DGK-alpha

Protein Families: Druggable Genome


Protein Pathways: Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways,

Phosphatidylinositol signaling system

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DGKA ([RC222395], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DGKA. Positive lysates [LY400535] (100ug) and [LC400535] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 5 different cell lines by using anti-DGKA monoclonal antibody (1:200).





Flow cytometric Analysis of Jurkat cells, using anti-DGKA antibody ([TA504050]), (Red), compared to a nonspecific negative control antibody, (Blue).