

Product datasheet for TA504077AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CCK4 (PTK7) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1G4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1G4

Applications: WB

Recommended Dilution: WB 1:1000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PTK7(NP_690619) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: protein tyrosine kinase 7 (inactive)

Database Link: NP 690619

Entrez Gene 71461 MouseEntrez Gene 5754 Human

Q13308

Background: Receptor protein tyrosine kinases transduce extracellular signals across the cell membrane. A

subgroup of these kinases lack detectable catalytic tyrosine kinase activity but retain roles in signal transduction. The protein encoded by this gene is a member of this subgroup of tyrosine kinases and may function as a cell adhesion molecule. This gene is thought to be expressed in colon carcinomas but not in normal colon, and therefore may be a marker for or may be involved in tumor progression. Four transcript variants encoding four different

isoforms have been found for this gene. [provided by RefSeq]






Synonyms: CCK-4; CCK4

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PTK7 ([RC223416], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PTK7.