

## Product datasheet for **TA504092AM**

### **MKRN1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2C8]**

#### **Product data:**


<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI2C8
<b>Applications:</b>	FC, IHC, WB
<b>Recommended Dilution:</b>	WB 1:1000, IHC 1:150, FLOW 1:100
<b>Reactivity:</b>	Human, Mouse
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG2b
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Full length human recombinant protein of human MKRN1(NP_038474) produced in HEK293T cell.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.5 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Conjugation:</b>	Biotin
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Predicted Protein Size:</b>	53.2 kDa
<b>Gene Name:</b>	makorin ring finger protein 1
<b>Database Link:</b>	<a href="#">NP_038474</a> <a href="#">Entrez Gene 54484 Mouse</a> <a href="#">Entrez Gene 23608 Human</a> <a href="#">Q9UHC7</a>
<b>Background:</b>	The Makorin ring finger protein-1 gene (MKRN1) is a highly transcribed, intron-containing source for a family of intronless mammalian genes encoding a novel class of zinc finger proteins. Phylogenetic analyses indicate that the MKRN1 gene is the ancestral founder of this gene family (Gray et al., 2000 [PubMed 10843807]). [supplied by OMIM, Mar 2008]
<b>Synonyms:</b>	RNF61




[View online »](#)

Protein Families: Druggable Genome


**Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MKRN1 (Cat# [RC206848], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MKRN1 (Cat# [TA504092]). Positive lysates [LY415580] (100ug) and [LC415580] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])







Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])




Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-MKRN1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA504092])



Flow cytometric Analysis of HeLa cells, using anti-MKRN1 antibody ([TA504092]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-MKRN1 antibody ([TA504092]), (Red), compared to a nonspecific negative control antibody, (Blue).