

Product datasheet for TA504138M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GIRK1 (KCNJ3) Mouse Monoclonal Antibody [Clone ID: OTI1C5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C5
Applications: IF, WB

Recommended Dilution: WB 1:500~2000, IF 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 279-501 of human

KCNJ3(NP_002230) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.81 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 56.4 kDa

Gene Name: potassium inwardly rectifying channel subfamily J member 3

Database Link: NP 002230

Entrez Gene 16519 MouseEntrez Gene 50599 RatEntrez Gene 488355 DogEntrez Gene 696473

MonkeyEntrez Gene 3760 Human

P48549






Background:

Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. The encoded protein, which has a greater tendency to allow potassium to flow into a cell rather than out of a cell, is controlled by G-proteins and plays an important role in regulating heartbeat. It associates with three other G-protein-activated potassium channels to form a heteromultimeric pore-forming complex. [provided by RefSeq]


Synonyms: GIRK1; KGA; KIR3.1

Protein Families: Druggable Genome, Ion Channels: Potassium, Transmembrane

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KCNJ3 ([RC205322], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KCNJ3. Positive lysates [LY400811] (100ug) and [LC400811] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-KCNJ3 monoclonal antibody.





Anti-KCNJ3 mouse monoclonal antibody ([TA504138]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY KCNJ3 ([RC205322]).