

## Product datasheet for **TA504155M**

### DGKA Mouse Monoclonal Antibody [Clone ID: OTI8G5]

#### Product data:

|                         |   |
|-------------------------|---|
| Product Type:           | Primary Antibodies  |
| Clone Name:             | OTI8G5  |
| Applications:           | IF, WB  |
| Recommended Dilution:   | WB 1:500~2000, IF 1:100   |
| Reactivity:             | Human, Dog, Monkey  |
| Host:                   | Mouse   |
| Isotype:                | IgG2b   |
| Clonality:              | Monoclonal  |
| Immunogen:              | Full length human recombinant protein of human DGKA(NP_001336) produced in HEK293T cell.  |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| Concentration:          | 1 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)   |
| Conjugation:            | Unconjugated  |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 82.5 kDa  |
| Gene Name:              | diacylglycerol kinase alpha   |
| Database Link:          | <a href="#">NP_001336</a><br><a href="#">Entrez Gene 100855691 Dog</a> <a href="#">Entrez Gene 710559 Monkey</a> <a href="#">Entrez Gene 1606 Human</a><br><a href="#">P23743</a>   |
| Background:             | The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It acts as a modulator that competes with protein kinase C for the second messenger diacylglycerol in intracellular signaling pathways. It also plays an important role in the resynthesis of phosphatidylinositols and phosphorylating diacylglycerol to phosphatidic acid. Alternative splicing occurs at this locus and four transcript variants encoding the same protein have been identified. [provided by RefSeq] |



[View online »](#)

**Synonyms:** DAGK; DAGK1; DGK-alpha


**Protein Families:** Druggable Genome

**Protein Pathways:** Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways, Phosphatidylinositol signaling system


### Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DGKA ([RC222395], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DGKA. Positive lysates [LY400535] (100ug) and [LC400535] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-DGKA monoclonal antibody.



Anti-DGKA mouse monoclonal antibody ([TA504155]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DGKA ([RC222395]).