

## **Product datasheet for TA504171S**

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

## IGJ (JCHAIN) Mouse Monoclonal Antibody [Clone ID: OTI2B1]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI2B1

**Applications:** FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IGJ(NP\_653247) produced in HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 17.9 kDa

**Gene Name:** joining chain of multimeric IgA and IgM

Database Link: NP 653247

Entrez Gene 3512 Human


P01591

Synonyms: IGCJ; IGJ; JCH







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IGJ ([RC207932], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IGJ. Positive lysates [LY403400] (100ug) and [LC403400] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-IGJ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504171])



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-IGJ mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504171])





Flow cytometric Analysis of Jurkat cells, using anti-IGJ antibody ([TA504171]), (Red), compared to a nonspecific negative control antibody, (Blue).