

Product datasheet for TA504173

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IGJ (JCHAIN) Mouse Monoclonal Antibody [Clone ID: OTI3B3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3B3

Applications: FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IGJ(NP_653247) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.83 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 17.9 kDa

Gene Name: joining chain of multimeric IgA and IgM

Database Link: NP 653247

Entrez Gene 3512 Human


P01591

Synonyms: IGCJ; IGJ; JCH






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IGJ (Cat# [RC207932], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IGJ(Cat# TA504173). Positive lysates [LY403400] (100ug) and [LC403400] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-IGJ mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-IGJ mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-IGJ monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).