

# **Product datasheet for TA504173AM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### IGI (ICHAIN) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3B3]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3B3

**Applications:** FC, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human IGJ(NP\_653247) produced in HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 17.9 kDa

**Gene Name:** joining chain of multimeric IgA and IgM

Database Link: NP 653247

Entrez Gene 3512 Human


P01591

Synonyms: IGCJ; IGJ; JCH







## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IGJ (Cat# [RC207932], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IGJ(Cat# [TA504173]). Positive lysates [LY403400] (100ug) and [LC403400] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-IGJ mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504173])



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-IGJ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504173])



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-IGJ monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).