

Product datasheet for TA504194S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

IGJ (JCHAIN) Mouse Monoclonal Antibody [Clone ID: OTI3H7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3H7
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IGJ(NP_653247) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.43 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 17.9 kDa

Gene Name: joining chain of multimeric IgA and IgM

Database Link: NP 653247


Entrez Gene 3512 Human

P01591


Synonyms: IGCJ; IGJ; JCH



Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IGJ ([RC207932], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IGJ. Positive lysates [LY403400] (100ug) and [LC403400] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-IGJ monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).







Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-IGJ mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504194])

Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-IGJ mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504194])