

Product datasheet for TA504249

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ALOX15 Mouse Monoclonal Antibody [Clone ID: OTI8C5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8C5
Applications: IHC, WB
Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse

Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ALOX15(NP_001131) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 74.6 kDa

Gene Name: arachidonate 15-lipoxygenase

Database Link: NP 001131

Entrez Gene 246 Human

P16050


Synonyms: 12-LOX; 15-LOX-1; 15LOX-1

Protein Families: Druggable Genome


Protein Pathways: Arachidonic acid metabolism, Linoleic acid metabolism, Metabolic pathways




Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALOX15 ([RC206621], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALOX15. Positive lysates [LY400456] (100ug) and [LC400456] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-ALOX15 mouse monoclonal antibody. (TA504249). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-ALOX15 mouse monoclonal antibody. (TA504249). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.