

## Product datasheet for **TA504257**

### Iduronate 2 sulfatase (IDS) Mouse Monoclonal Antibody [Clone ID: OTI1D9]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI1D9
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human IDS(NP_000193) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	59.2 kDa
Gene Name:	iduronate 2-sulfatase
Database Link:	<a href="#">NP_000193</a> <a href="#">Entrez Gene 15931 Mouse</a> <a href="#">Entrez Gene 3423 Human</a> <a href="#">P22304</a>

**Background:** Iduronate-2-sulfatase is required for the lysosomal degradation of heparan sulfate and dermatan sulfate. Mutations in this X-chromosome gene that result in enzymatic deficiency lead to the sex-linked Mucopolysaccharidosis Type II, also known as Hunter Syndrome. Iduronate-2-sulfatase has a strong sequence similarity with human arylsulfatases A, B, and C, and human glucosamine-6-sulfatase. Multiple alternatively spliced transcript variants that encode different protein isoforms have been described. [provided by RefSeq]



[View online »](#)

**Synonyms:** MPS2; SIDS  
**Protein Families:** Druggable Genome  
**Protein Pathways:** Glycosaminoglycan degradation, Lysosome, Metabolic pathways


### Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IDS ([RC219187], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IDS. Positive lysates [LY424863] (100ug) and [LC424863] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-IDS monoclonal antibody.




Western blot analysis of extracts (10ug) from a mouse cell line by using anti-IDS monoclonal antibody (1:200).




Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-IDS monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.







Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-IDS mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Anti-IDS mouse monoclonal antibody (TA504257) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IDS ([RC219187]).



HEK293T cells transfected with either [RC219187] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-IDS antibody (TA504257), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-IDS antibody (TA504257), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-IDS antibody (TA504257), (Red), compared to a nonspecific negative control antibody, (Blue).