

## Product datasheet for **TA504281S**

### **RUVBL2 Mouse Monoclonal Antibody [Clone ID: OTI1A6]**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	OTI1A6
<b>Applications:</b>	FC, IF, IHC, WB
<b>Recommended Dilution:</b>	WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100
<b>Reactivity:</b>	Human, Dog, Rat, Monkey, Mouse
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Human recombinant protein fragment corresponding to amino acids 113-370 of human RUVBL2(NP_006657) produced in E.coli.
<b>Formulation:</b>	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
<b>Concentration:</b>	0.6 mg/ml
<b>Purification:</b>	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
<b>Conjugation:</b>	Unconjugated
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Predicted Protein Size:</b>	51 kDa
<b>Gene Name:</b>	RuvB like AAA ATPase 2
<b>Database Link:</b>	<a href="#">NP_006657</a> <a href="#">Entrez Gene 20174 Mouse</a> <a href="#">Entrez Gene 292907 Rat</a> <a href="#">Entrez Gene 476418 Dog</a> <a href="#">Entrez Gene 100428794 Monkey</a> <a href="#">Entrez Gene 10856 Human</a> <a href="#">Q9Y230</a>



[View online »](#)

**Background:**


This gene encodes the second human homologue of the bacterial RuvB gene. Bacterial RuvB protein is a DNA helicase essential for homologous recombination and DNA double-strand break repair. Functional analysis showed that this gene product has both ATPase and DNA helicase activities. This gene is physically linked to the CGB/LHB gene cluster on chromosome 19q13.3, and is very close (55 nt) to the LHB gene, in the opposite orientation. [provided by RefSeq]

**Synonyms:**


CGI-46; ECP51; INO80J; REPTIN; RVB2; TIH2; TIP48; TIP49B

**Protein Families:**


Transcription Factors

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RUVBL2 (Cat# [RC200933], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RUVBL2(Cat# [TA504281]). Positive lysates [LY401993] (100ug) and [LC401993] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-RUVBL2 monoclonal antibody.




Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-RUVBL2 monoclonal antibody (1:200).




Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-RUVBL2 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon).




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-RUVBL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504281])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-RUVBL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504281])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-RUVBL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504281])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-RUVBL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504281])



Anti-RUVBL2 mouse monoclonal antibody ([TA504281]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY RUVBL2 ([RC200933]).



HEK293T cells transfected with either [RC200933] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-RUVBL2 antibody ([TA504281]), and then analyzed by flow cytometry.