

Product datasheet for TA504402

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NLN Mouse Monoclonal Antibody [Clone ID: OTI3G8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3G8

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NLN(NP_065777) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1.1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80.5 kDa

Gene Name: neurolysin

Database Link: NP 065777

Entrez Gene 75805 MouseEntrez Gene 117041 RatEntrez Gene 696884 MonkeyEntrez Gene

<u>57486 Human</u>

Q9BYT8

Background: This gene encodes a member of the metallopeptidase M3 protein family that cleaves

neurotensin at the Pro10-Tyr11 bond, leading to the formation of neurotensin(1-10) and neurotensin(11-13). The encoded protein is likely involved in the termination of the neurotensinergic signal in the central nervous system and in the gastrointestinal tract.





Synonyms: AGTBP; EP24.16; MEP; MOP

Protein Families: Druggable Genome, Protease

Protein Pathways: Renin-angiotensin system

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NLN ([RC212447], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NLN. Positive lysates [LY412353] (100ug) and [LC412353] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NLN monoclonal antibody.