

Product datasheet for **TA504411BM**


PDRG1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4F12]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI4F12
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:150, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDRG1(NP_110442) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	15.3 kDa
Gene Name:	p53 and DNA damage regulated 1
Database Link:	NP_110442 Entrez Gene 68559 Mouse Entrez Gene 296278 Rat Entrez Gene 81572 Human Q9NUG6
Synonyms:	C20orf126; PDRG




[View online »](#)

Product images:


HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PDRG1 (Cat# [RC203661], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDRG1 (Cat# [TA504411]). Positive lysates [LY410685] (100ug) and [LC410685] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-PDRG1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA504411])



Flow cytometric Analysis of HeLa cells, using anti-PDRG1 antibody ([TA504411]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-PDRG1 antibody ([TA504411]), (Red), compared to a nonspecific negative control antibody, (Blue).