

Product datasheet for TA504423BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SPG7 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1D2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1D2

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 300-573 of human

SPG7(NP_003110) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 88.1 kDa

Gene Name: SPG7 matrix AAA peptidase subunit, paraplegin

Database Link: NP 003110

Entrez Gene 234847 MouseEntrez Gene 353231 RatEntrez Gene 6687 Human

O9UO90






Background:

This gene encodes a nuclear-encoded mitochondrial metalloprotease protein that is a member of the AAA (ATPases associated with a variety of cellular activities) protein family. Members of this protein family share an ATPase domain and have roles in diverse cellular processes including membrane trafficking, intracellular motility, organelle biogenesis, protein folding, and proteolysis. Two transcript variants encoding distinct isoforms have been identified for this gene. Mutations associated with this gene cause autosomal recessive spastic paraplegia 7. [provided by RefSeq, Jul 2008]

Synonyms: CAR; CMAR; PGN; SPG5C

Protein Families: Protease, Transmembrane

Product images:



Negative control E. coli lysate (Left lane) or E. coli lysate containing recombinant protein fragment for human SPG7 (NP_003110) gene (amino acids 300-573) (Right lane). Equivalent amounts (5 ug per lane) were separated by SDS-PAGE and then immunoblotted with anti-SPG7.