

Product datasheet for TA504458S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PALS1 Mouse Monoclonal Antibody [Clone ID: OTI5D10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5D10

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MPP5(NP_071919) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 77.1 kDa

Gene Name: protein associated with LIN7 1, MAGUK family member

Database Link: NP 071919

Entrez Gene 314259 RatEntrez Gene 64398 Human

O8N3R9



Background:


Members of the peripheral membrane-associated guanylate kinase (MAGUK) family function in tumor suppression and receptor clustering by forming multiprotein complexes containing distinct sets of transmembrane, cytoskeletal, and cytoplasmic signaling proteins. All MAGUKs contain a PDZ-SH3-GUK core and are divided into 4 subfamilies, DLG-like (see DLG1; MIM 601014), ZO1-like (see TJP1; MIM 601009), p55-like (see MPP1; MIM 305360), and LIN2-like (see CASK; MIM 300172), based on their size and the presence of additional domains (Tseng et al., 2001 [PubMed 11311936]). MPP5 is a member of the p55-like MAGUK subfamily. [supplied by OMIM]


Synonyms: PALS1

Protein Families: Druggable Genome

Protein Pathways: Tight junction

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MPP5 ([RC224752], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MPP5. Positive lysates [LY411659] (100ug) and [LC411659] (20ug) can be purchased separately from OriGene.

Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and MPP5-Knockout HeLa cells (KO, Cat# [LC834761]) were separated by SDS-PAGE and immunoblotted with anti-MPP5 monoclonal antibody [TA504458] (1:200). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.