

Product datasheet for **TA504553AM**

ATIC Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1D2]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D2
Applications:	FC, WB
Recommended Dilution:	WB 1:500~2000, FLOW 1:100
Reactivity:	Human, Dog, Rat, Monkey, Mouse
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ATIC(NP_004035) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	64.4 kDa
Gene Name:	5-aminoimidazole-4-carboxamide ribonucleotide formyltransferase/IMP cyclohydrolase
Database Link:	NP_004035 Entrez Gene 81643 Rat Entrez Gene 108147 Mouse Entrez Gene 488513 Dog Entrez Gene 694819 Monkey Entrez Gene 471 Human P31939
Background:	This gene encodes a bifunctional protein that catalyzes the last two steps of the de novo purine biosynthetic pathway. The N-terminal domain has phosphoribosylaminoimidazolecarboxamide formyltransferase activity, and the C-terminal domain has IMP cyclohydrolase activity. A mutation in this gene results in AICA-ribosiduria. [provided by RefSeq]




[View online »](#)

Synonyms: AICAR; AICARFT; HEL-S-70p; IMPCHASE; PURH


Protein Families: Stem cell - Pluripotency

Protein Pathways: Metabolic pathways, One carbon pool by folate, Purine metabolism


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ATIC (Cat# [RC203490], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ATIC(Cat# [TA504553]). Positive lysates [LY418255] (100ug) and [LC418255] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ATIC monoclonal antibody.



HEK293T cells transfected with either [RC203490] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ATIC antibody ([TA504553]), and then analyzed by flow cytometry.