

Product datasheet for TA504570BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ATIC Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4A10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A10

Applications: WB

Recommended Dilution: WB 1:500~1000

Reactivity: Human, Dog, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ATIC(NP_004035) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 64.4 kDa

Gene Name: 5-aminoimidazole-4-carboxamide ribonucleotide formyltransferase/IMP cyclohydrolase

Database Link: NP 004035

Entrez Gene 81643 RatEntrez Gene 108147 MouseEntrez Gene 488513 DogEntrez Gene 471

<u>Human</u> <u>P31939</u>

Background: This gene encodes a bifunctional protein that catalyzes the last two steps of the de novo

purine biosynthetic pathway. The N-terminal domain has

phosphoribosylaminoimidazolecarboxamide formyltransferase activity, and the C-terminal domain has IMP cyclohydrolase activity. A mutation in this gene results in AICA-ribosiduria.

[provided by RefSeq]






Synonyms: AICAR; AICARFT; HEL-S-70p; IMPCHASE; PURH


Protein Families: Stem cell - Pluripotency

Protein Pathways: Metabolic pathways, One carbon pool by folate, Purine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ATIC ([RC203490], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ATIC. Positive lysates [LY418255] (100ug) and [LC418255] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-ATIC monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).