

# **Product datasheet for TA504744S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NQO2 Mouse Monoclonal Antibody [Clone ID: OTI3C11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3C11

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Dog, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NQO2(NP\_000895) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 25.7 kDa

**Gene Name:** N-ribosyldihydronicotinamide:quinone reductase 2

Database Link: NP 000895

Entrez Gene 606932 DogEntrez Gene 707675 MonkeyEntrez Gene 4835 Human

P16083

**Background:** NQO2 (EC 1.10.99.2) is a flavoprotein that catalyzes the 2-electron reduction of various

quinones, redox dyes, and the vitamin K menadione. NQO2 predominantly uses

dihydronicotinamide riboside (NRH) as the electron donor (summary by Wu et al., 1997


[PubMed 9367528]). [supplied by OMIM, Jul 2010]

Synonyms: DHQV; DIA6; NMOR2; QR2







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NQO2 ([RC202889], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NQO2. Positive lysates [LY424463] (100ug) and [LC424463] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-NQO2 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).



Western blot analysis of extracts (10ug) from a mouse tissue by using anti-NQO2 monoclonal antibody (1:200).