

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA504748BM

DBT Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1G2]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	TI1G2	
Applications:	FC, IF, WB	
Recommended Dilution:	:2000, FLOW 1:100	
Reactivity:	man, Mouse, Rat	
Host:	se	
lsotype:	1	
Clonality:	noclonal	
Immunogen:	Full length human recombinant protein of human DBT(NP_001909) produced in HEK293T cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.	
Concentration:	0.5 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	HRP	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	46.4 kDa	
Gene Name:	dihydrolipoamide branched chain transacylase E2	
Database Link:	<u>NP_001909</u> <u>Entrez Gene 13171 MouseEntrez Gene 29611 RatEntrez Gene 1629 Human</u> <u>P11182</u>	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	DBT Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1G2] – TA504748BM			
Background:	The branched-chain alpha-keto acid dehydrogenase complex (BCKD) is an inner- mitochondrial enzyme complex involved in the breakdown of the branched-chain amino acids isoleucine, leucine, and valine. The BCKD complex is thought to be composed of a core of 24 transacylase (E2) subunits, and associated decarboxylase (E1), dehydrogenase (E3), and regulatory subunits. This gene encodes the transacylase (E2) subunit. Mutations in this gene result in maple syrup urine disease, type 2. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq]			
Synonyms:	BCATE2; BCKAD-E2; BCKADE2; BCOADC-E2; E2; E2B			
Protein Families	: Druggable Genome			
Protein Pathway	/s: Metabolic pathways, Valine, leucine and isoleucine degradation			

Product images:


170	_	
130	-	
100	-	
70	-	
55	-	
40	-	
35	-	
25	-	
15	-	
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DBT (Cat# [RC201998], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DBT(Cat# [TA504748]). Positive lysates [LY419653] (100ug) and [LC419653] (20ug) can be purchased separately from OriGene.



Anti-DBT mouse monoclonal antibody ([TA504748]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY DBT ([RC201998]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



HEK293T cells transfected with either [RC201998] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DBT antibody ([TA504748]), and then analyzed by flow cytometry.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US